

**UNCLASSIFIED**

---

**AD 274 000**

*Reproduced  
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA**



---

**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

AFOSR2167


274 000

(2)

274 000

AD No.

ASTIA FILE COPY



FILE COPY

Return to

ASTIA

RLINGTON HALL STATION

RLINGTON 12, VIRGINIA

Attn: TIRS

DEPARTMENT OF PHYSICS

THE HEBREW UNIVERSITY

JERUSALEM, ISRAEL

# 1.6 0

ASTIA  
RECEIVED  
APR 11 1962  
TISIA A

Monitoring Agency No.

Astia Document No.

**PARAMAGNETIC RESONANCE SPECTRA OF RARE EARTH IONS  
IN THE CRYSTAL FIELD OF CALCIUM FLUORIDE <sup>2</sup>**

**W. Low  
Department of Physics  
The Hebrew University  
Jerusalem, Israel**

**Technical Note No. 18**

**January 1962**


**Contract No. AF 61(052)-59**


**Presented at the International Conference on Magnetism and Crystallography,  
Kyoto, Japan, in September, 1961.**

**■**

**The research in this document has been sponsored in part by the Air Force  
Office of Scientific Research, U.S.A.F., through its European Office of  
Aerospace Research.**

ABSTRACT

  
Paramagnetic resonance data of various rare earth ions  
in the crystal field of ~~CaF<sub>2</sub>~~ <sup>Calcium Fluoride</sup> are presented. It is found  
that there are at least three types of symmetries, cubic,  
axial and trigonal. <sup>Some</sup> In particular the spectra of ~~Co<sup>3+</sup>, V<sup>3+</sup>~~ <sup>Co<sup>3+</sup>, V<sup>3+</sup></sup>,  
~~Er<sup>3+</sup>, Nd<sup>3+</sup>~~ <sup>Er<sup>3+</sup>, Nd<sup>3+</sup></sup> are presented.

  
—  
\* CERIUM (3+), YTTERBIUM (3+), ERBIUM (3+), AND NEODYMIUM (3+)

We have studied the paramagnetic resonance spectra of rare earth ions in the crystal field of calcium fluoride. These spectra can be classified as belonging to three types: a. spectra which indicate that the rare earth ions are exposed to a cubic field; b. spectra in which the anisotropic behaviour suggests an axial crystal field along the cubic directions; c. spectra in which the axial distortion is along the body diagonals of the cube.

The calcium ion with fluorite structure is surrounded by a cube of eight nearest fluorine ions. The trivalent rare earth ions substitute for the calcium ion. However, a charge compensation has to take place to preserve the neutrality of the crystal. From measurements of the X-ray lattice constants<sup>1</sup> and of the ionic conductivity<sup>2</sup> it has been shown that in well annealed crystals the main disorder introduced into the crystal by the addition of the rare earth ion is the formation of  $F^-$  interstitial (see Fig. 1). The empty sites which are available for the interstitial  $F^-$  ion are located at the center of the adjacent cube along one of the cubic axes. Supporting evidence for the  $F^-$  interstitial formation is found in the work of Bleaney et al.<sup>3</sup>

With proper thermal treatment, part or a large fraction of the interstitial fluorine ions can be removed from the adjacent cube next to the rare earth ion.<sup>4</sup> In this case, the point symmetry of the rare earth ions approximates  $O_h$ . In some crystals, however, the predominant axial distortion is found from paramagnetic resonance spectra to be along the body diagonals of the cube. The optical spectra shows a number of strong bands and very poor ultraviolet transmission. It is likely that one of the fluorine ions at the corner of the cube is replaced by  $O^{2-}$  or  $OH^-$  ion. Supporting evidence is found that on heating the crystal in air this spectrum increases in intensity.

The respective crystal fields are given by

$$a) V \text{ cubic} = B_4^0 \left[ O_4^0 + 5(O_4^4 + O_4^{-4}) \right] + B_6^0 \left[ O_6^0 - 21(O_6^4 + O_6^{-4}) \right]$$

$$b) V \text{ axial} = B_2^0 O_2^0 + B_4^0 O_4^0 + B_6^0 O_6^0 + B_4^4 (O_4^4 + O_4^{-4}) + B_6^4 (O_6^4 + O_6^{-4})$$

$$c) V \text{ diagonal} = B_2^0 O_2^0 + B_4^0 O_4^0 + B_6^0 O_6^0 + B_4^3 (O_4^3 + O_4^{-3}) + B_6^3 (O_6^3 + O_6^{-3})$$

where  $B_n^m$  are crystal field coefficients proportional to  $\frac{r^n}{R^{n+1}}$  and  $O_n^m$  are

operators transforming the corresponding spherical Legendre polynomials.

The magnetic properties of these three types of spectra are very different and we shall illustrate this in a few selected examples. A full report of the magnetic properties of the rare earth ions in calcium fluoride will appear elsewhere.

$$\text{Er}^{3+} : {}^4\text{H}_{15/2}$$

a) In the cubic field of calcium fluoride, the 16-fold degeneracy splits into 3 quarters ( $\Gamma_8$ ) and two doublets. The lowest levels are given by

$$-258.86 \text{ c} - 162 \text{ d}, \Gamma_8$$

$$-26 \text{ c} - 312 \text{ d}, \Gamma_7$$

where c and d are parameters signifying the fourth and sixth order contribution. For a point charge model they are given by

$$c = \frac{2^3}{3^3 \cdot 11.13} \frac{Z_0^2}{R^5} \quad d = \frac{2^4}{3^3} \frac{10}{11.13^2} \frac{Z_0^2}{R^7}$$

The observed g factor is  $6.785 - 0.002$ . This is consistent with a  $\Gamma_7$  ground state which gives g calculated = 6.79. A lowest  $\Gamma_7$  level indicates that the rare earth spectrum is dominated to a large extent by the sixth order term in the potential.

b) An axial spectrum has been observed by Baker et al<sup>6</sup> and by the author with  $g_{\parallel} = 7.76$  and  $g_{\perp} = 6.253$ . This is consistent with the same doublet level except now exposed to an axial field since  $\frac{g_{\parallel} + 2g_{\perp}}{3} \sim g$  cubic

c) A set of three lines, arising from the three inequivalent sites with axial symmetry along the three body diagonal directions, is observed in some crystals.<sup>7</sup> Each ion can be described by  $g$  factors

$$g_{\parallel} = 2.23, \quad g_{\perp} = 8.9 \text{ and again } \frac{g_{\parallel} + 2g_{\perp}}{3} \sim g \text{ cubic.}$$

The  $g$  factors can be explained to arise from a doublet with

$$a | \pm 13/2 \rangle + b | \pm 7/2 \rangle + c | \pm 1/2 \rangle + d | \mp 5/2 \rangle + e | \mp 11/2 \rangle$$

with the coefficients  $b$  and  $d$  larger than  $a$ ,  $c$  or  $e$ .

$$\text{Yb}^{3+} : {}^2F_{7/2}$$

(a) The cubic field suggests that  $\Gamma_7$  is the ground state level. The wave function of the doublet is given by

$$\sqrt{\frac{3}{4}} | \pm 5/2 \rangle - \sqrt{\frac{1}{4}} | \mp 3/2 \rangle$$

and the calculated  $g$  factor =  $24/7$ . The observed  $g$  factor of  $3.426 \pm 0.001$  is in essential agreement.<sup>8</sup>

(c) The spectrum along the body diagonals can be represented by the spin Hamiltonian

$$H = g_{\parallel} \beta H_z S_z + g_{\perp} \beta [H_x S_x + H_y S_y] + A S_z I_z + B(S_x I_x + S_y I_y) + P(I_z^2 - 1/3 I(I+1))$$

with  $S = \frac{1}{2}$  and  $I = \frac{1}{2}$  or  $I = 5/2$ .

The observed  $g$  factor is  $g_{\parallel} = 1.323 \pm 0.001$  and  $g_{\perp} = 4.387 \pm 0.002$  showing that this is the  $\Gamma_7$  anisotropic level. This Kramer's doublet can be approximated with

$$0.940 | \pm 1/2 \rangle + 0.264 | \pm 7/2 \rangle + 0.209 | \mp 5/2 \rangle$$



The hyperfine part is expressed as

$$A^{171} = 354.6$$

$$B^{171} = 1168$$

$$A^{173} = 96$$

$$B^{173} = 320$$

$$P = 85 \pm 3. \quad \frac{g_n}{g_J} \frac{B}{A} \sim 1$$

all in  $10^{-4}$  cm. These results indicate a large quadrupole moment of about 2.4 barns.

$$Nd^{3+} \quad 4I_{9/2}$$

(a) Calculations show that the ground state for all ratios of the fourth to sixth order parameters is given by a quartet  $\sqrt{8}$ . The wave functions and g factors are given by

<u>g value</u>	<u>transition</u>	<u>relative intensity</u>
2.24	$0.8722 -5/2\rangle - 0.4892 +3/2\rangle \rightarrow 0.8722 +5/2\rangle - 0.4892 -3/2\rangle$	20
1.15	$0.8722 +5/2\rangle - 0.4892 +3/2\rangle \rightarrow 0.05410 +9/2\rangle - 0.2981 +1/2\rangle + 0.9530 +7/2\rangle$	4.4
4.54	$0.5410 +9/2\rangle - 0.2981 +1/2\rangle + 0.9530 -7/2\rangle - 0.05410 -9/2\rangle - 0.2981 -1/2\rangle + 0.9530 +7/2\rangle$	0.075

The spin Hamiltonian of the  $\sqrt{8}$  level in a cubic field is outlined by Bleaney<sup>9</sup> applies here. The spectrum has been investigated and corresponds approximately to this Hamiltonian.

$$H = gB(H_i S_i) + fB(H_i S_i^3) \quad i = x, y, z.$$

The observed and calculated g factors are given in Table I<sup>10</sup>.

(b) The axial field spectrum has been investigated by Bleaney et al<sup>3</sup> and remeasured by us<sup>10</sup>. The g factors are given  $g_n = 4.412$  and  $g_J = 1.301$ . The ground state can be described possibly by

$$= a|+9/2\rangle + b|+1/2\rangle + c|+7/2\rangle$$

(c) A weak spectrum is also observed with the axial field along the body diagonals.

We have, in addition, observed the cubic field spectrum of  $\text{Ce}^{3+}$ ,  $\text{Dy}^{3+}$  and  $\text{Gd}^{3+}$ , the axial field spectra of these ions. The axial spectra of ions with even number of electrons such as  $\text{Tb}^{3+}$ , and  $\text{Tm}^{3+}$  has been observed as well.<sup>7</sup>

A large part of this work has been performed in collaboration with Mr. M. Dvir, U. Rosenberger and Dr. G. Vincow.

---

TABLE I

Measured and calculated g factor of  $\Gamma_8$  ground state of  $\text{Nd}^{3+}$ .

Direction	g exp.	g calc.
100	2.26	2.24
	1.10	1.15
110	1.45	1.46
	1.6	1.7


REFERENCES





1. E. Zintl and A. Udgard, Z. anorg. u. allgem. Chemie 240, 150 (1939)
  2. R.W. Ure, J. Chem. Phys. 26, 1363 (1957)
  3. B. Bleaney, P.M. Llewellyn, D.W. Jones, Proc. Phys. Soc. B (London), 69, 858 (1956)
  4. E. Friedman and W. Low, J. Chem. Phys. 33, 1275 (1960)
  5. M. Dvir and W. Low, Proc. Phys. Soc. (London) 74, 136 (1960)
  6. J.M. Baker, W. Hayes and D.W. Jones, Proc. Phys. Soc. (London) 73, 943 (1959).
  7. W. Low and U. Rosenberger (to be published).
  8. W. Low, Phys. Rev. 118, 1608 (1960)
  9. B. Bleaney, Proc. Phys. Soc. (London) 73, 937 (1959)
  10. G. Vincow and W. Low, Phys. Rev. 122, 1390 (1961).
-

Fig. 1.

Crystal Structure of  $\text{CaF}_2$

The figure indicates the possible defect sites near the rare earth ion substituting for the calcium ion.



-   $\text{F}^-$
-   $\text{Ca}^{2+}$  or  $\text{X}^{3+}$
-   $\text{F}^-$  interstitial
-   $\text{O}^{--}$  or  $\text{HO}^-$

THE HEBREW UNIVERSITY  
JERUSALEM, ISRAEL  
DEPARTMENT OF PHYSICS  
TECHNICAL REPORT NO. 18  
JANUARY 1962

AF 61(052)-59  
TN-18  
PHYSICS

PARAMAGNETIC RESONANCE SPECTRA OF RARE EARTH IONS IN THE  
CRYSTAL FIELD OF CALCIUM FLUORIDE

W. LOW

ABSTRACT: Paramagnetic resonance data of various rare earth ions in the crystal field of  $\text{CaF}_2$  is presented. It is found that there are at least three types of symmetries, cubic, axial and trigonal. In particular the spectra of  $\text{Ce}^{3+}$ ,  $\text{Yb}^{3+}$ ,  $\text{Er}^{3+}$ ,  $\text{Nd}^{3+}$  are presented.

THE HEBREW UNIVERSITY  
JERUSALEM, ISRAEL  
DEPARTMENT OF PHYSICS  
TECHNICAL REPORT NO. 18  
JANUARY 1962

AF 61(052)-59  
TN-18  
PHYSICS

PARAMAGNETIC RESONANCE SPECTRA OF RARE EARTH IONS IN THE  
CRYSTAL FIELD OF CALCIUM FLUORIDE

W. LOW

ABSTRACT: Paramagnetic resonance data of various rare earth ions in the crystal field of  $\text{CaF}_2$  is presented. It is found that there are at least three types of symmetries, cubic, axial and trigonal. In particular the spectra of  $\text{Ce}^{3+}$ ,  $\text{Yb}^{3+}$ ,  $\text{Er}^{3+}$ ,  $\text{Nd}^{3+}$  are presented.

THE HEBREW UNIVERSITY  
JERUSALEM, ISRAEL  
DEPARTMENT OF PHYSICS  
TECHNICAL REPORT NO. 18  
JANUARY 1962

AF 61(052)  
TN-18  
PHYSICS

PARAMAGNETIC RESONANCE SPECTRA OF RARE EARTH IONS IN THE  
CRYSTAL FIELD OF CALCIUM FLUORIDE

W. LOW

ABSTRACT: Paramagnetic resonance data of various rare earth ions in the crystal field of  $\text{CaF}_2$  is presented. It is found that there are at least three types of symmetries, cubic, axial and trigonal. In particular the spectra of  $\text{Ce}^{3+}$ ,  $\text{Yb}^{3+}$ ,  $\text{Er}^{3+}$ ,  $\text{Nd}^{3+}$  are presented.

THE HEBREW UNIVERSITY  
JERUSALEM, ISRAEL  
DEPARTMENT OF PHYSICS  
TECHNICAL REPORT NO. 18  
JANUARY 1962

AF 61(052)-59  
TN-18  
PHYSICS

PARAMAGNETIC RESONANCE SPECTRA OF RARE EARTH IONS IN THE  
CRYSTAL FIELD OF CALCIUM FLUORIDE

W. LOW

ABSTRACT: Paramagnetic resonance data of various rare earth ions in the crystal field of  $\text{CaF}_2$  is presented. It is found that there are at least three types of symmetries, cubic, axial and trigonal. In particular the spectra of  $\text{Ce}^{3+}$ ,  $\text{Yb}^{3+}$ ,  $\text{Er}^{3+}$ ,  $\text{Nd}^{3+}$  are presented.